2920X

Teklad Irradiated Global Soy Protein-Free Extruded Rodent Diet

Product Description- 2920X is a fixed formula, irradiated, extruded diet manufactured with high quality ingredients designed to support gestation, lactation, and growth of rodents. 2920X does not contain alfalfa or soybean meal, thus minimizing the occurrence of phytoestrogens. Typical isoflavone concentrations (daidzein + genistein agylcone equivalents) range from non-detectable to 20 mg/kg. Exclusion of alfalfa reduces chlorophyll, improving optical imaging clarity. Absence of fish meal minimizes the presence of nitrosamines. Diet is exposed to irradiation dose not exceeding 50 kGy. **Related codes 2020X, 2020SX (autoclavable).**

Macronutrients Crude Protein % 19.4 Fat (acid hydrolysis) a % 6.5 Carbohydrate (available) b % 47.0 Crude Fiber % 2.7 Neutral Detergent Fiber c % 12.3 Ash % 5.1 Energy Density d kcal/g (kJ/g) 3.1 (13.0) Calories from Protein % 24 Calories from Protein % 16 Calories from Carbohydrate % 60 Minerals Calcium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4 Magnesium % 0.2
Fat (acid hydrolysis) a % 6.5 Carbohydrate (available) b % 47.0 Crude Fiber % 2.7 Neutral Detergent Fiber c % 12.3 Ash % 5.1 Energy Density d kcal/g (kJ/g) 3.1 (13.0) Calories from Protein % 24 Calories from Fat % 16 Calories from Carbohydrate % 60 Winerals Calcium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Carbohydrate (available) b % 47.0 Crude Fiber % 2.7 Neutral Detergent Fiber c % 12.3 Ash % 5.1 Energy Density d kcal/g (kJ/g) 3.1 (13.0) Calories from Protein % 24 Calories from Fat % 16 Calories from Carbohydrate % 60 Minerals Solium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Crude Fiber % 2.7 Neutral Detergent Fiber c % 12.3 Ash % 5.1 Energy Density d kcal/g (kl/g) 3.1 (13.0) Calories from Protein % 24 Calories from Fat % 16 Calories from Carbohydrate % 60 Minerals Calcium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Neutral Detergent Fiber c % 12.3 Ash % 5.1 Energy Density d kcal/g (kJ/g) 3.1 (13.0) Calories from Protein % 24 Calories from Fat % 16 Calories from Carbohydrate % 60 Minerals O.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Ash % 5.1 Energy Density ^d kcal/g (kl/g) 3.1 (13.0) Calories from Protein % 24 Calories from Fat % 16 Calories from Carbohydrate % 60 Minerals Calcium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Energy Density ^d kcal/g (kJ/g) 3.1 (13.0) Calories from Protein % 24 Calories from Fat % 16 Calories from Carbohydrate % 60 Minerals Calcium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Calories from Protein % 24 Calories from Fat % 16 Calories from Carbohydrate % 60 Minerals Calcium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Calories from Protein % 24 Calories from Fat % 16 Calories from Carbohydrate % 60 Minerals Calcium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Calories from Carbohydrate % 60 Minerals Second of the control of the contro
Minerals Calcium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Calcium % 0.9 Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Phosphorus % 0.7 Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Non-Phytate Phosphorus % 0.4 Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Sodium % 0.1 Potassium % 0.4 Chloride % 0.4
Potassium % 0.4 Chloride % 0.4
Chloride % 0.4
Magnesium % 0.2
1410E11C310111 /0 U.Z
Zinc mg/kg 60
Manganese mg/kg 80
Copper mg/kg 15
lodine mg/kg 6
Iron mg/kg 200
Selenium mg/kg 0.23
Amino Acids
Aspartic Acid % 1.1
Glutamic Acid % 3.5
Alanine % 1.2
Glycine % 0.7
Threonine % 0.6
Proline % 1.9
Serine % 0.9
Leucine % 2.3
Isoleucine % 0.7
Valine % 0.9
Phenylalanine % 1.0
Tyrosine % 0.5
Methionine % 0.5
Cystine % 0.3
Lysine % 0.9
Histidine % 0.4
Arginine % 0.8
Tryptophan % 0.2



Ingredients (in descending order of inclusion)- Ground wheat, ground corn, corn gluten meal, wheat middlings, soybean oil, calcium carbonate, dicalcium phosphate, brewers dried yeast, L-lysine, iodized salt, magnesium oxide, choline chloride, DL-methionine, calcium propionate, L-tryptophan, vitamin E acetate, menadione sodium bisulfite complex (source of vitamin K activity), manganous oxide, ferrous sulfate, zinc oxide, niacin, calcium pantothenate, copper sulfate, pyridoxine hydrochloride, riboflavin, thiamin mononitrate, vitamin A acetate, calcium iodate, vitamin B_{12} supplement, folic acid, biotin, vitamin D_3 supplement, cobalt carbonate.

Standard	Product F	orm· F	vtrudod

010110101		
Vitamins		
Vitamin A ^{e, f}	IU/g	15.0
Vitamin D ₃ ^{e, g}	IU/g	1.5
Vitamin E	IU/kg	110
Vitamin K ₃ (menadione)	mg/kg	50
Vitamin B ₁ (thiamin)	mg/kg	17
Vitamin B ₂ (riboflavin)	mg/kg	15
Niacin (nicotinic acid)	mg/kg	75
Vitamin B ₆ (pyridoxine)	mg/kg	18
Pantothenic Acid	mg/kg	33
Vitamin B ₁₂ (cyanocobalamin)	mg/kg	0.08
Biotin	mg/kg	0.40
Folate	mg/kg	4
Choline	mg/kg	1200
Fatty Acids		
C16:0 Palmitic	%	0.6
C18:0 Stearic	%	0.1
C18:1ω9 Oleic	%	1.1
C18:2ω6 Linoleic	%	2.6
C18:3ω3 Linolenic	%	0.3
Total Saturated	%	0.8
Total Monounsaturated	%	1.1
Total Polyunsaturated	%	2.9
Other		
Cholesterol	mg/kg	

Shelf life: With proper storage, diet is suitable for use out to 9 months.

www.inotivco.com/shelf-life-of-diets-used-in-research

For nutrients not listed, insufficient data is available to quantify.

Nutrient data represent the best information available, calculated from published values and direct analytical testing of raw materials and finished product. Nutrient values may vary due to the natural variations in the ingredients, analysis, and effects of processing.



^a Ether extract is used to measure fat in pelleted diets, while an acid hydrolysis method is required to recover fat in extruded diets. Compared to ether extract, the fat value for acid hydrolysis will be approximately 1% point higher.

^b Carbohydrate (available) is calculated by subtracting neutral detergent fiber from total carbohydrates.

^c Neutral detergent fiber is an estimate of insoluble fiber, including cellulose, hemicellulose, and lignin. Crude fiber methodology underestimates total fiber.

^d Energy density is a calculated estimate of *metabolizable energy* based on the Atwater factors assigning 4 kcal/g to protein, 9 kcal/g to fat, and 4 kcal/g to available carbohydrate.

^e Indicates added amount but does not account for contribution from other ingredients.

f 1 IU vitamin A = 0.3 μg retinol

g 1 IU vitamin D = 25 ng cholecalciferol