## inotiv

Product Description- 7922 is a fixed and open formula, irradiated diet

manufactured with high quality ingredients and designed to support growth and reproduction of rodents. Typical isoflavone concentrations (daidzein + genistein aglycone equivalents) range from 270 to 400 mg/kg. Diet is exposed to irradiation dose not exceeding 50 kGy. Related code 7022.

Ingredients (in descending order of inclusion)- Ground corn, ground wheat, dehulled soybean meal, fish meal, wheat middlings, dried skimmed milk, dehydrated alfalfa meal, corn gluten meal, soybean oil, brewers dried yeast, cane molasses, dicalcium phosphate, calcium carbonate, salt, choline chloride, ferrous sulfate, manganous oxide, vitamin E acetate, vitamin  $\mathrm{B}_{\mathrm{12}}$  supplement, menadione sodium bisulfite complex (source of vitamin K activity), calcium pantothenate, niacin, zinc oxide, copper sulfate, riboflavin, biotin, thiamin mononitrate, vitamin D<sub>3</sub> supplement, vitamin A acetate, pyridoxine hydrochloride, folic acid, calcium iodate, cobalt carbonate.

Standard Product Form: Pellet

Macronutrients		
Crude Protein	%	22.5
Fat (ether extract) <sup>a</sup>	%	5.2
Carbohydrate (available) <sup>b</sup>	%	42.7
Crude Fiber	%	3.7
Neutral Detergent Fiber <sup>c</sup>	%	11.7
Ash	%	6.7
Energy Density <sup>d</sup>	kcal/g (kJ/g)	3.1 (13.0)
Calories from Protein	%	29
Calories from Fat	%	15
Calories from Carbohydrate	%	56
Minerals		
Calcium	%	1.2
Phosphorus	%	1.0
Non-Phytate Phosphorus	%	0.7
Sodium	%	0.3
Potassium	%	0.9
Chloride	%	0.6
Magnesium	%	0.2
Zinc	mg/kg	56
Manganese	mg/kg	105
Copper	mg/kg	13
Iodine	mg/kg	2
Iron	mg/kg	350
Selenium	mg/kg	0.31
Amino Acids		
Aspartic Acid	%	1.9
Glutamic Acid	%	3.6
Alanine	%	1.4
Glycine	%	1.3
Threonine	%	0.9
Proline	%	1.6
Serine	%	1.3
Leucine	%	1.9
Isoleucine	%	1.0
Valine	%	1.2
Phenylalanine	%	1.1
Tyrosine	%	0.9
Methionine	%	0.5
Cystine	%	0.3
Lysine	%	1.3
Histidine	%	0.6
Arginine	%	1.4

Vitamin A <sup>e, f</sup>	IU/g	6.1
Vitamin D <sub>3</sub> <sup>e, g</sup>	IU/g	5.1
Vitamin E	IU/kg	50
Vitamin K <sub>3</sub> (menadione)	mg/kg	4
Vitamin B <sub>1</sub> (thiamin)	mg/kg	16
Vitamin B <sub>2</sub> (riboflavin)	mg/kg	6
Niacin (nicotinic acid)	mg/kg	78
Vitamin B <sub>6</sub> (pyridoxine)	mg/kg	11
Pantothenic Acid	mg/kg	31
Vitamin B <sub>12</sub> (cyanocobalamin)	mg/kg	0.07
Biotin	mg/kg	0.33
Folate	mg/kg	3
Choline	mg/kg	1830
Fatty Acids		
C16:0 Palmitic	%	0.8
C18:0 Stearic	%	0.2
C18:1ω9 Oleic	%	1.1
C18:2ω6 Linoleic	%	2.2
C18:3ω3 Linolenic	%	0.2
Total Saturated	%	1.1
Total Monounsaturated	%	1.3
Total Polyunsaturated	%	2.4
Other		
Cholesterol	mg/kg	60

## www.inotivco.com/shelf-life-of-diets-used-in-research

<sup>a</sup> Ether extract is used to measure fat in pelleted diets, while an acid hydrolysis method is required to recover fat in extruded diets. Compared to ether extract, the fat value for acid hydrolysis will be approximately 1% point higher.

- <sup>b</sup> Carbohydrate (available) is calculated by subtracting neutral detergent fiber from total carbohydrates.
- <sup>c</sup> Neutral detergent fiber is an estimate of insoluble fiber, including cellulose, hemicellulose, and lignin. Crude fiber methodology underestimates total fiber.
- <sup>d</sup> Energy density is a calculated estimate of *metabolizable energy* based on the Atwater factors assigning 4 kcal/g to protein, 9 kcal/g to fat, and 4 kcal/g to available carbohvdrate.
- <sup>e</sup> Indicates added amount but does not account for contribution from other ingredients.
- <sup>f</sup> 1 IU vitamin A = 0.3 μg retinol

Vitamins

<sup>g</sup> 1 IU vitamin D = 25 ng cholecalciferol

For nutrients not listed, insufficient data is available to quantify.

Nutrient data represent the best information available, calculated from published values and direct analytical testing of raw materials and finished product. Nutrient values may vary due to the natural variations in the ingredients, analysis, and effects of processing.

Tryptophan



0.3

%